아밀로이드 병리 및 영상

예 병 석

연세의대 신경과학교실

Amyloid pathology & imaging

Byoung Seok Ye, MD, PhD

Department of Neurology Yonsei University College of Medicine

ORIGINAL ARTICLES

Imaging Brain Amyloid in Alzheimer's Disease with Pittsburgh Compound-B

William E. Klunk, MD, PhD,¹ Henry Engler, MD,² Agneta Nordberg, MD, PhD,^{3,4} Yanming Wang, PhD,⁵ Gunnar Blomqvist, PhD,² Daniel P. Holt, BS,⁵ Mats Bergström, PhD,² Irina Savitcheva, MD,² Guo-feng Huang, PhD,⁵ Sergio Estrada, PhD,⁵ Birgitta Aulin, MSc,⁴ Manik L. Debnath, MS,¹ Julien Barletta, BS,⁵ Julie C. Price, PhD,⁵ Johan Sandell, PhD,² Brian J. Lopresti, BS,⁵ Anders Wall, PhD,² Pernilla Koivisto, PhD,² Gunnar Antoni, PhD,² Chester A. Mathis, PhD,² and Bengt Långström, PhD^{2,6}

Klunk WE, 2004, *Ann Neurol*

Various tracers for β -amyloid

Thioflavin T

Pittsburgh Compound B

Flutemetamol

Florbetaben

Florbetapir

Radioactive analogues of thioflavin T (Fluorescent dye for amyloid staining)

¹¹C PiB had 20 min half-life

¹⁸F tracers have longer half-life

Filippi L, 2018, *Am J Nucl Med Mol Imaging*

Florbetaben PET imaging to detect amyloid beta plaques in Alzheimer disease: Phase 3 study

Osama Sabri^{1,2,3}, Marwan N. Sabbagh^{1,4}, John Seibyl¹, Henryk Barthel¹, Hiroyasu Akatsu^{1,5,6}, Yasuomi Ouchi¹, Kohei Senda¹, Shigeo Murayama¹, Kenji Ishii¹, Masaki Takao^{1,6}, Thomas G. Beach⁷, Christopher C. Rowe⁸, James B. Leverenz^{9,10}, Bernardino Ghetti¹¹, James W. Ironside⁶, Ana M. Catafau¹, Andrew W. Stephens⁶, Andre Mueller¹², Norman Koglin¹³, Anja Hoffmann¹⁴, Katrin Roth¹⁵, Cornelia Reininger¹⁶, Walter J. Schulz-Schaeffer¹, and for the Florbetaben Phase 3 Study Group²

Sabri O, 2015, *Alzheimer's & Dementia*

Performance of [¹⁸F]flutemetamol amyloid imaging against the neuritic plaque component of CERAD and the current (2012) NIA-AA recommendations for the neuropathologic diagnosis of Alzheimer's disease

Neuroimaging

From 76 modified CERAD $A\beta$ -positive and 30 $A\beta$ -negative cases

Only visual rating done, No quantitative analyses performed

Accuracy

1) Visual rating: 90.6%

-Sensitivity: 69/76 = 90.8%

-Specificity: 27/30 = 90.0%

Salloway S, 2017, *Alzheimer's & Dementia DADM*

Cerebral PET with florbetapir compared with neuropathology at autopsy for detection of neuritic amyloid- β plaques: a prospective cohort study

Christopher M Clark¹, Michael J Prineas², Thomas G Beach³, Barry J Bredt⁴, R Edward Coleman⁵, P Murali Doraiswamy⁶, Adam S Finkelstein⁷, Eric M Reiman⁸, Marwan N Sabbagh⁹, Carl H Siderowky¹⁰, Julie A Schneider¹¹, Anupa Avona¹², Alan P Carpenter¹³, Matthew L Fitzee¹⁴, Abhinav D Joshi¹⁵, Michael J Knudsen¹⁶, Ming Lu¹⁷, Mark A Mintzer¹⁸, Daniel M Skovronsky¹⁹, for the A4-45-A16 Study Group²⁰

Clark CM, 2012, *Lancet Neurol*

Clinical implication of amyloid PET

- ◆ Imaging biomarker supported diagnoses of AD
 - ◆ AD dementia, MCI d/t AD, SCD d/t AD, and preclinical AD
 - ◆ Exclusion of AD from diagnosis (special caution needed)
- ◆ Detection of mixed amyloid pathology in other type of dementia
- ◆ Diagnosis of cerebral amyloid angiopathy

Imaging biomarker supported diagnosis of AD

Imaging biomarker supported diagnosis of AD

Regional SUVR	Left	Right
Caudal Middle Frontal	3.17	3.06
Lateral Frontal Opercularis	2.05	2.01
Supernasal Gyrus	2.19	2.17
Lateral Orbitofrontal	3.14	3.20
Lateral Frontal Orbitals	1.80	1.92
Middle Temporal	2.19	2.20
Medial Orbitofrontal	3.17	3.22
Rostrol Middle Frontal	2.27	2.20
Rostrol Anterior Cingulate	2.23	2.30
Inferior Cingulate Gyrus	2.10	2.05
Superior Parietal	2.32	2.18
Lateral Frontal Opercularis	2.06	1.98
Caudal Anterior Cingulate	2.17	2.20
Superior Frontal Gyrus	3.17	3.13
Precentral	2.44	2.30
Inferior Parietal	2.27	2.24
Posterior Cingulate	2.26	2.23
Superior Temporal	2.22	2.01

Global FBB
SUVR = 2.18

Exclusion of AD from diagnosis (special caution needed)

Exclusion of AD from diagnosis (special caution needed)

Regional SUVR	Left	Right
Caudal Middle Frontal	1.23	1.21
Lateral Frontal Opercularis	1.18	1.22
Supernasal Gyrus	1.18	1.18
Lateral Orbitofrontal	1.21	1.25
Lateral Frontal Orbitals	1.16	1.22
Middle Temporal	1.21	1.22
Medial Orbitofrontal	1.11	1.18
Rostrol Middle Frontal	1.19	1.14
Rostrol Anterior Cingulate	1.18	1.20
Inferior Cingulate Gyrus	1.23	1.27
Superior Parietal	1.24	1.24
Lateral Frontal Opercularis	1.13	1.18
Caudal Anterior Cingulate	1.24	1.22
Superior Frontal Gyrus	1.16	1.16
Precentral	1.19	1.20
Inferior Parietal	1.21	1.26
Posterior Cingulate	1.22	1.23
Superior Temporal	1.24	1.22

Global FBB
SUVR = 1.20

Exclusion of AD from diagnosis (special caution needed)

Borderline case

Exclusion of AD from diagnosis (special caution needed)

Regional positive case

Cerebral amyloid angiopathy

Florbetapir-PET to diagnose cerebral amyloid angiopathy

A prospective study

Abstract

Objective: We hypothesized that florbetapir, a Food and Drug Administration-approved PET tracer, could distinguish cerebral amyloid angiopathy (CAA)-related intracerebral hemorrhage (ICH) from hippocampal CAH (CAH).

Methods: We prospectively enrolled survivors of primary ICH related to probable CAA per Boston Criteria (n = 23) and HTN-ICH (n = 18) without dementia. All patients underwent florbetapir-PET and multimodal MRI, and patients with CAA had additional Pittsburgh compound B (PiB) PET. Amyloid burden was assessed quantitatively (standard uptake value ratio [SUVR]) and visually classified as positive or negative.

Results: The CAA and HTN-ICH groups had similar ages (65.9 vs 67.1, sex, and leukocyte volume fraction [LV] vs 20.4%, all p > 0.16). Florbetapir uptake and PiB retention strongly correlated in patients with CAA both globally (cortical SUVR $r = 0.86$, $p < 0.001$) and regionally in lower cortical (all $r > 0.6$, all $p < 0.01$). Mean global cortical florbetapir uptake was 4.2-fold higher in CAA than HTN-ICH (SUVR 2.42 ± 0.27 vs 1.15 ± 0.08, $p < 0.001$), as was mean occipital SUVR (2.44 ± 0.12 vs 1.17 ± 0.08, $p < 0.001$), even after correcting for global SUVR ($p < 0.01$). Visual ratings for occipital florbetapir demonstrated inter-PET rater agreement $k = 0.77$ (range positive for all 12 tracers) vs HTN-ICH $k = 0.27$ (range 0 to 100% non-PiB) (Cohen's kappa, 100% non-PiB) (95% CI).

Conclusions: Florbetapir appears to label vascular amyloid in patients with CAA-related ICH. The approved florbetapir brain visual rating method can have diagnostic value in appropriate clinical settings.

Classification of evidence: This study provides Class II evidence that florbetapir-PET provides a sensitivity of 100%, 95% confidence interval (CI) 65%-100% and specificity of 89% (95% CI 65%-99%) for determination of probable CAA among cognitively normal patients. <https://doi.org/10.1212/000000000000000000>

Gurol ME, 2016, *Neurology*

Summary + Take home message

- ◆ Amyloid PET is a useful imaging biomarker for exact diagnosis of AD
 - ◆ Special caution is needed to rule out AD
 - ◆ Regional positivity is common especially in cases with mixed pathologies (LBD, Vascular, etc)
- ◆ Amyloid PET is a biomarker for amyloid, not for AD (amyloid + tau)
 - ◆ AD, LB variant of AD, DLB with amyloid, AD with DLB, and even normal cognition subjects can show amyloid deposition
 - ◆ Biomarkers for disease staging or differential diagnosis are needed
 - ◆ AD and LBD could have direct interaction through occipital amyloid deposition
 - ◆ CAA could be diagnosed with amyloid PET