Update on Neuromuscular Disorders

석흥열
계명대학교

Hung Youl Seok, MD

Department of Neurology Keimyung University, Dongsan Medical Center, Daegu, Korea

1

Myasthenia gravis

Satisfactory Response With Achieving Maintenance Low-Dose Prednisone in Generalized Myasthenia Gravis

J Clin Neuromusc Dis 2018;20:49-59

	Whole Cohort			Not Taking an SSI			Taking an SSI		
	At 2 years, N = 81	At 4 years, N = 56	At 6 years, N = 32	At 2 years, N = 45	At 4 years, N = 34*	At 6 years, n = 17†	At 2 years, N = 36	At 4 years, N = 22*	At 6 years, N = 15
SR% n/N; if LDP ≤ 7.5 mg	46.9 (38/81)	57.1 (32/56)	62.5 (20/32)	41.4 (20/45)	61.7 (22/34)	58.8 (10/17)	50.0 (18/36)	45.4 (10/22)	66.7 (10/15)
Sensitivity analysis									
SR% (n/N; if LDP ≤ 10 mg)	58.0 (47/81)	69.6 (39/56)	68.7 (22/32)	55.6 (25/45)	71.0 (24/34)	70.5 (12/17)	61.1 (22/36)	68.1 (15/22)	66.7 (10/15)
SR% (n/N; if LDP ≤ 5 mg)	43.2 (35/81)	57.1 (32/56)	18/32	56.3 (19/45)	61.7 (22/34)	58.8 (10/17)	41.4 (16/36)	45.4 (10/22)	66.7 (8/15)
SR% (n/N) maintained for 1 y	43.2 (35/81)	53.6 (30/56)	19/32	40.0 (18/45)	61.8 (21/34)	52.9 (9/17)	47.2 (17/36)	40.9 (9/22)	66.7 (10/15)
SR% (n/N) if "not taking an SSI" group was defined as no exposure to an SSI at all	—	—	—	51.8 (17/31)	64.3 (18/28)	56.3 (9/16)	50.0 (18/36)	45.4 (10/22)	66.7 (10/15)
SR% (n/N) if "not taking an SSI" group was defined as <3 months exposure to SSI	—	—	—	52.8 (19/36)	61.3 (18/28)	56.3 (9/16)	50.0 (18/36)	45.4 (10/22)	66.7 (10/15)
Secondary outcomes									
Remission/minimal manifestation (%; n/N)	87.7 (71/81)	89.3 (50/56)	100 (32/32)	86.7 (39/45)	91.1 (32/34)	100 (17/17)	88.9 (32/36)	81.8 (18/22)	100 (15/15)
No moderate or severe worsening (%; n/N)	97.5 (79/81)	92.8 (52/56)	100 (31/32)	97.8 (44/45)	94.1 (32/34)	100 (17/17)	97.2 (35/36)	90.9 (20/22)	93.3 (14/15)

Safety and efficacy of eculizumab in anti-acetylcholine receptor antibody-positive refractory generalised myasthenia gravis (REGAIN): a phase 3, randomised, double-blind, placebo-controlled, multicentre study

James F Howard¹, Kimiaki Utsugisawa, Michael Benatar, Hitoyuki Mura, Richard J Barohn, Isabel Ila, Saiji Jacob, John Vissing, Ted M Burns, John T Kissel, Srikanth Muppala, Richard J Nowak, Fanny O'Brien, Jing-jing Wang, Renato Monteggia, in collaboration with the REGAIN Study Group*

Lancet Neurol 2017; 16: 976-86

	Eculizumab (n=62)	Placebo (n=63)	Difference (95% CI)	p value*
Prespecified worst-rank ANCOVA score†				
MG-ADL‡	56.6 (4.5)	68.3 (4.5)	-11.7 (-24.3 to 0.96)	0.0698
QMG	54.7 (4.5)	70.7 (4.5)	-16.0 (-28.5 to -3.4)	0.0129
MGC	57.3 (4.5)	67.7 (4.5)	-10.5 (-23.1 to 2.1)	0.1026
MG-QOL15	55.5 (4.6)	69.7 (4.5)	-14.3 (-27.0 to -1.6)	0.0281
Prespecified sensitivity repeated-measures model analysis with immunosuppressive treatments as covariates§				
MG-ADL	-4.1 (0.5)	-2.3 (0.5)	-1.8 (-3.2 to -0.5)	0.0077
QMG	-4.6 (0.6)	-1.7 (0.6)	-2.9 (-4.6 to -1.2)	0.0007
MGC	-7.9 (1.0)	-4.6 (1.0)	-3.3 (-5.9 to -0.6)	0.0168
MG-QOL15	-13.8 (1.6)	-6.7 (1.6)	-7.1 (-11.3 to -3.0)	0.0009
Post-hoc sensitivity worst-rank ANCOVA score¶				
MG-ADL	54.8 (4.5)	70.2 (4.4)	-15.4 (-27.8 to -2.9)	0.0160
QMG	53.9 (4.5)	71.6 (4.4)	-17.7 (-30.1 to -5.3)	0.0055
MGC	56.1 (4.5)	69.0 (4.4)	-12.9 (-25.4 to -0.5)	0.0414
MG-QOL15	54.6 (4.5)	70.6 (4.5)	-16.0 (-28.6 to -3.4)	0.0134

Event	Number of events	Number of patients experiencing an event (%)	Event rate (events per 100 patient-years)
Most common adverse events* (≥ 10% of all patients, N = 117)			
Headache	71	44 (37.6)	31.3
Nasopharyngitis	76	37 (31.6)	33.5
Diarrhea	40	27 (23.1)	17.6
Upper respiratory tract infection	55	27 (23.1)	24.2
Myasthenia gravis†	40	23 (19.7)	17.6
Arthralgia	29	22 (18.8)	12.8
Nausea	26	21 (17.9)	11.5
Pain in extremity	21	18 (15.4)	9.3
Cough	21	17 (14.5)	9.3
Fatigue	21	17 (14.5)	9.3
Urinary tract infection	32	17 (14.5)	14.1
Influenza	24	16 (13.7)	10.6
Gastroenteritis	15	14 (12.0)	6.6
Bronchitis	22	13 (11.1)	9.7
Pyrexia	17	15 (13.1)	7.5
Fall	24	12 (10.3)	10.6
Deaths	3	3 (2.6)	1.3

Characteristics of patients with remission vs. patients without remission at last follow-up

Variables	Total (n=56)	Remission at last follow-up (n=24)	No remission at last follow-up (n=32)	p value
Baseline characteristics before RTX start				
Female gender	34 (60.7)	14 (58.3)	20 (62.5)	0.75
Age at diagnosis of MG in years, median (IQR)	41.5 (24.3; 65.8)	58 (33; 71)	29.5 (20; 57.5)	0.01
Age at RTX start in years, median (IQR)	47.5 (33; 71)	67 (40; 72)	40.5 (31; 64.3)	0.029
Duration of diagnosed MG before RTX start in years, median (IQR)	4 (1.3; 10.8)	2.5 (1; 6.8)	6 (2; 12)	0.09
Antibody type				
Anti-AchR-ab	39 (69.6)	14 (58.3)	25 (78.1)	0.11
Anti-MuSK-ab	14 (25.0)	10 (41.7)	4 (12.5)	0.013
Double seronegative	3 (5.4)	0	3 (9.4)	
Thymoma	8 (14.3)	4 (16.7)	4 (12.5)	0.71
Number of steroid-sparing therapies before RTX start, mean (SD)	2.9 (1.4)	2.2 (1.1)	3.4 (1.4)	0.001
High-cost therapies (PLEX/IA or IVIG) before RTX start	47 (83.9)	17 (70.8)	30 (93.8)	0.03
Disease severity before RTX start according to MGFA-CC				
Mild disease (MGFA-CC < 3)	15 (26.8)	10 (41.7)	5 (15.6)	0.029
Moderate to severe disease (MGFA-CC ≥ 3b)	33 (58.9)	12 (50)	21 (65.6)	0.24

	Prednisone group	Thymectomy plus prednisone group	Estimated difference (95% CI)	p value
Time-weighted mean alternate-day prednisone dose, mg	49.0 (29.2; 33)	25.9 (20.7; 35)	23.1 (10.9 to 35.2)	0.0003*
Peralised time-weighted mean alternate-day prednisone dose, mg				
Method 1†	66.2 (36.7; 33)	31.0 (31.8; 35)	35.2 (18.6 to 51.9)	<0.0001*
Method 2‡	60.6 (34.6; 33)	28.3 (27.9; 35)	32.3 (17.2 to 47.5)	<0.0001*
Time-weighted and time-specific mean Myasthenia Gravis Activities of Daily Living score§				
Month 0-60	3.26 (2.77; 32)	1.61 (1.46; 34)	1.65 (0.54 to 2.75)	0.0044*
Month 48	2.55 (3.02; 29)	1.10 (1.51; 31)	1.45 (0.20 to 2.71)	0.0245*
Month 60	2.04 (2.63; 24)	1.23 (1.75; 26)	0.81 (-0.48 to 2.07)	0.21*
Azathioprine use	19/33 (58%)	7/35 (20%)	37.6% (16.1 to 59.0%)	0.0014*
Plasma exchange use	4/33 (12%)	5/35 (14%)	-2.1% (-18.2 to 13.9)	0.73*
Intravenous immunoglobulin use	11/33 (33%)	3/35 (9%)	24.8% (6.2 to 43.3)	0.0162*
Minimal Manifestation Status				
Month 48	15/29 (52%)	23/31 (74%)	-22.5% (-46.3 to 1.4)	0.07*
Month 60	14/24 (58%)	23/26 (88%)	-30.1% (-53.4 to 6.9)	0.0236*
MG-QOL15				
Month 39	13.1 (14.0; 32)	4.8 (9.2; 33)	8.2 (2.3 to 14.1)	0.0029**
Month 48	9.0 (10.1; 29)	4.9 (7.9; 30)	4.1 (0.6 to 8.8)	0.13**
Month 60	7.7 (9.24; 24)	7.8 (10.9; 26)	-0.1 (-5.9 to 5.6)	0.96**

Who will benefit from thymectomy for myasthenia gravis? Is there any role for this procedure in elderly patients?

Ryo Otsuka, Kazuhiro Ueda, Toshiki Tanaka, Junichi Murakami, Masataro Hayashi, Kimikazu Hamano
Ann Transl Med 2019;7(1):4

Variables	Surgical outcome		P
	Improved	Not improved	
Age (years)	49.5±18.9	63.0±10.9	0.0683
Age (≥65/<65 years)	4/18	5/3	0.0192
Gender (male/female)	6/16	3/5	0.59
Smoking history (yes/no)	5/17	2/6	0.90
PS (0/1/2/3)	12/7/1/2	4/2/1/1	0.86
BMI (kg/m ²)	22.4±3.5	21.7±3.1	0.64
VC (% of predicted)	98.4±15.3	99.9±31.1	0.86
Thymoma (present/absent)	10/12	1/7	0.0976
Autoantibody (elevated/normal)	19/3	8/0	0.27
MGFA (I/II/III/IV/V)	4/13/3/1/1	1/5/2/0/0	0.43
Onset (<50/≥50 years)	8/14	2/6	0.56
Length from onset to surgery (<1/≥1 year)	17/5	2/6	0.0086
QMG score	12.2±4.8	13.3±5.1	0.61
Surgical approach (open/VATS)	2/20	2/6	0.26

Results of a multiple logistic regression analysis for the risk of treatment failure

Variables	Reference	Odds ratio	P
Length from onset to surgery	≥1 year	15.6	0.0237
Age at surgery	≥65 years	12.0	0.0405

Emergence of Myasthenia Gravis with Myositis in a Patient Treated with Pembrolizumab for Thymic Cancer

So-Young Huh
J Clin Neurol 2018;14(1):115-117

Case reports of neuromuscular disorder after pembrolizumab administration

Age/sex reference	Underlying disease	Symptom onset* (weeks)	Symptoms	Diagnosis	CK (IU/L)	ACHR ab*	Immuno-therapy	Outcome
34/F	Thymic cancer (our case)	8	EOM limitation, ptosis, neck weakness, dyspnea, dysphagia, and hypophonia	MG with myositis	2,125	+	IVIg, IVMP, PLEX, prednisone	Partial recovery
86/F ²	Metastatic melanoma	6	EOM limitation, ptosis, proximal limb weakness, fatigue, and dysphonia	Necrotic myositis	1,499	-	IVMP, PLEX	Recovery
69/F ²	Metastatic melanoma	9	EOM limitation, ptosis, dyspnea, and general weakness	MG	↑	-	IVMP, PLEX	No significant recovery, died due to brain metastasis
75/M ⁴	Metastatic melanoma, well-controlled MG	5	Ptosis, dyspnea, and neck and proximal limb weakness	Exacerbation of MG	NR	+	IVIg, prednisone	Recovery
59/F ²	Metastatic melanoma, well-controlled MG	12	Dyspnea, dysphagia with dysarthria, and neck and proximal limb weakness	Exacerbation of MG	NR	-	PLEX, IVIg, prednisone	Recovery
71/F ²	Metastatic melanoma	12	EOM limitation, ptosis, dysphagia, dysarthria, and neck and proximal limb weakness	MG	1,200	-	Prednisone	Recovery, died due to underlying cancer
78/M ⁴	Metastatic melanoma	8	Ptosis, facial weakness, dyspnea, dysphagia, proximal limb weakness, and myalgia	Necrotic myositis over a NMJ disorder	1,284	-	PLEX, prednisone	No significant recovery, passed away
63/M ⁴	Metastatic melanoma	2	EOM limitation, ptosis, facial weakness, dyspnea, and periorbital edema with erythema	MG with myositis	10,386	+	IVIg, IVMP, PLEX, prednisone	No significant recovery, died
85/F ²	Metastatic melanoma	4.5	Diplopia and ptosis	Ocular MG	NR	-	IVIg, prednisone	Recovery

Inflammatory myopathy with myasthenia gravis

Thymoma association and polymyositis pathology
Naohiro Uchio, MD
Neural Neuroimmunol Neuroinflamm 2019;6:e535.

Patient	1	2	3	4	5	6	7	8	9	10
Age at IM/MG diagnosis (y)	59/50	73/56	53/52	50/38	51/43	49/36	41/52	78/78	70/70	60/60
Sex	Male	Female	Male	Female	Male	Female	Female	Male	Female	Female
MG diagnosis before IM diagnosis (y)	+9	+17	<1	+12	+14	+13	-5	<1	<1	<1
Thymoma	+	+	+	+	+	+	+	-	-	-
Disease activity of MG										
Ocular symptoms/ fluctuating weakness at IM diagnosis	-/+	+/-	+/-	+/-	+/-	+/-	+/-	+/-	+/-	+/-
Incremental responses on RNS at IM diagnosis	-	-	-	+	-	-	-	+	+	+
Positive edrophonium infusion test at IM diagnosis	NE	NE	NE	NE	NE	NE	-	+	NE	+
Anti-AChR Ab at MG-IM diagnosis (IU/mL)	ND/22	11/540	100/ND	80/300	16/7	41/10	ND/12	47.7	6.5	↓
CK level at MG diagnosis (IU/L)	ND	ND	231	ND	185	41	ND	263	1,140	282

Patient	1	2	3	4	5	6	7	8	9	10
Clinical features of IM at IM diagnosis										
History of transiently elevated CK levels	+	-	-	-	-	-	-	-	-	-
Progression of weakness at IM diagnosis	Rapid	No	Rapid	Chronic	Chronic	Rapid	Chronic	Rapid	Rapid	Chronic
Limb weakness	Severe	No-mild	Severe	Moderate	No-mild	Severe	No-mild	No-mild	No-mild	Moderate
Dropped head	+	-	+	-	+	-	-	+	-	-
Dysphagia	+	-	+	+	+	-	-	-	ND	-
Respiratory failure	+(NIPPV)	-	+(IM)	-	-	+(IM)	+(IM)	+(IM)	-	-
Cardiac involvement	-	-	-	Atrioseptal hypokinesia	-	VT and diffuse hypokinesia	-	-	VT and diffuse hypokinesia	-
Laboratory findings at IM diagnosis										
CK level (IU/L)	10,226	376	1,008	341	1,059	5,102	7,315	263	1,140	212
Myositis-specific autoantibodies	-	-	-	-	-	-	-	-	-	NE
Anti-titin Ab	-	NE	+	+	+	+	+	+	+	NE
Spontaneous activities at rest on rEMG	+	+	+	+	+	+	+	+	NE	-
Histopathologic findings										
PD-1 ⁺ cell infiltration	+	+	+	+	+	+	NE	+	+	+
PD-L1 upregulation on fibers ^a	++	-	+	-	+	+	NE	+	++	+
CTLA-4 ⁺ cell infiltration	+	-	+	+	-	-	NE	-	-	-
Classification based on ENMC criteria	Definite PM	Definite PM	Definite PM	Probable PM	Probable PM	Probable PM	Definite PM	Probable PM	Probable PM	Definite PM

Guillain-Barre Syndrome

Regional variation of Guillain-Barré syndrome

Alex Y. Doets,^{1,*} Christine Verboon,^{1,*} Bianca van den Berg,^{1,*} Thomas Harbo,² David R. Cornblath,³ Hugh J. Willison,⁴ Zhahirul Islam,⁵ Shahram Attarian,⁶ Fabio A. Barroso,⁷ Kathleen Bateman,⁸ Luana Benedetti,⁹ Peter van den Bergh,¹⁰ Carlos Casanovas,¹¹ Guido Cavaletti,¹² Govindsinh Chavada,⁴ Kristi G. Claeys,^{13,14} Efthimios Dardiotis,¹⁵ Amy Davidson,⁴ Pieter A. van Doorn,¹ Tom E. Feasby,¹⁶ Giuliana Galassi,¹⁷ Kenneth C. Gorson,¹⁸ Hans-Peter Hartung,¹⁹ Sung-Tsang Hsieh,²⁰ Richard A.C. Hughes,²¹ Isabel Illa,²² Badrul Islam,⁵ Susumu Kusonoki,²³ Satoshi Kùwabara,²⁴ Helmar C. Lehmann,²⁵ James A.L. Miller,²⁶ Quazi Deen Mohammad,²⁷ Soledad Monges,²⁸ Eduardo Nobile-Orazio,²⁹ Julio Pardo,³⁰ Yann Persson,³¹ Simon Rimadi,³² Luis Querol,³³ Stephen W. Reddel,³⁴ Ricardo C. Reisin,³⁴ Nortina Shahrizaila,³⁵ Soren H. Sindrup,¹⁶ Waheed Waqar,³⁷ Bart C. Jacobs^{3,8} and the IGOS Consortium[#]

Brain. 2018 Oct 1;141(10):2866-2877.

	Region			P-value
	Europe/Americas (n = 715)	Asia (n = 69)	Bangladesh (n = 125)	
Demographics				
Age, years (IQR)	55 (37-67)	50 (34-60)	28 (16-40)	<0.001
Male:female ratio (%)	418/297 (1.41)	42/27 (1.56)	84/41 (2.05)	0.18
Clinical features at entry				
MRC sum score, possible range 0-60 ^a (IQR)	48 (38-56)	49 (40-58)	22 (7-37)	<0.001
Sensory deficits (%)	463/686 (65)	37/68 (54)	35/120 (28)	<0.001
Cranial nerve involvement (%)	330/712 (46)	44/69 (64)	84/125 (67)	<0.001
Oculomotor weakness	106/712 (15)	26/69 (38)	51/125 (4)	<0.001
Facial weakness	220/712 (31)	28/69 (41)	32/125 (26)	0.10
Bulbar weakness	142/712 (20)	23/69 (33)	64/125 (51)	<0.001
Autonomic dysfunction (%)	189/714 (27)	7/69 (10)	28/125 (22)	0.01
Pain (%)	415/713 (58)	8/69 (12)	77/125 (62)	<0.001
Time from onset of weakness to admission, days (IQR)	3 (2-6)	4 (2-6)	4 (2-8)	0.01
Clinical features at nadir				
MRC sum score, possible range 0-60 ^a (IQR)	46 (30-54)	48 (34-58)	16 (3-32)	<0.001
GBS disability score (%)				
Unable to walk independently (>2)	478/626 (76)	50/66 (76)	100/107 (93)	<0.001
Sensory deficits (%)	408/508 (80)	37/63 (59)	29/100 (29)	<0.001
Cranial nerve involvement (%)	304/620 (49)	44/65 (68)	73/107 (68)	<0.001
Oculomotor weakness	84/620 (14)	25/65 (39)	51/107 (5)	<0.001
Facial weakness	220/620 (36)	31/65 (48)	32/107 (30)	0.06
Bulbar weakness	136/620 (22)	24/65 (37)	57/107 (53)	<0.001
Autonomic dysfunction (%)	184/626 (29)	11/66 (17)	30/107 (28)	0.09
Pain (%)	354/625 (57)	11/66 (17)	67/107 (63)	<0.001
Ventilator dependency (%)	121/715 (17)	17/69 (25)	36/125 (29)	0.004

	Region			
	Europe/Americas (n=715)	Asia (n=69)	Bangladesh (n=125)	P-value
Electrophysiology classification (%)				
Demyelinating	312/573 (55)	29/65 (45)	38/94 (40)	0.02
Axonal	335/73 (6)	4/65 (6)	34/94 (36)	<0.001
Inexcitable	105/73 (2)	1/65 (2)	9/94 (10)	<0.001
Equivoal	182/573 (32)	20/65 (31)	12/94 (10)	0.001
Normal	365/73 (6)	11/65 (17)	1/94 (1)	<0.001
Initial treatment (%)				
None	54/715 (7)	9/69 (13)	108/125 (86)	<0.001
IVIg	612/715 (86)	50/69 (73)	7/125 (6)	<0.001
PE	43/715 (6)	10/69 (15)	9/125 (7)	0.03
Other ^a	6/715 (1)	0/69 (0)	1/125 (1)	0.75
Time from onset of weakness to treatment, days (IQR)	4 (2-7)	5 (3-7)	7 (5-12)	0.003
Outcome				
Median time to independent walking, days (IQR)	63 (28-186)	39 (17-94)	95 (36-190)	0.002
Able to walk independently at 6 months (%)	331/418 (79)	36/41 (88)	60/97 (62)	<0.001
Able to walk independently at 12 months (%)	334/404 (83)	31/34 (91)	67/97 (69)	0.003
Mortality				
Patients deceased at 12 months (%)	23/486 (5)	1/45 (2)	19/114 (17)	<0.001

The electrodiagnosis of Guillain-Barré syndrome subtypes: Where do we stand?

Clinical Neurophysiology 129 (2018) 2586-2593

Hadden's criteria	Rajabally's criteria
1) Primary demyelinating At least one of the following in each of two nerves, or at least two of the following in one nerve if all others inexcitable and distal CMAP $\geq 10\%$ LLN • MCV < 90% LLN (85% if dCMAP < 50% LLN) • DML > 110% ULN (120% if dCMAP < 100% LLN) • pCMAP/dCMAP amplitude ratio < 0.5 and distal CMAP $\geq 20\%$ LLN • F-response latency > 120% ULN 2) Primary axonal • None of the above features of demyelination in any nerve (except one demyelinating feature allowed in one nerve if distal CMAP < 10% LLN) AND • dCMAP < 80% LLN in at least two nerves 3) Inexcitable • dCMAP absent in all nerves (or present in only one nerve with distal CMAP < 10% LLN) 4) Equivoal • Does not exactly fit criteria for any other group	1) Acute inflammatory demyelinating polyneuropathy (AIDP) At least one of the following in at least two nerves: • MCV < 70% LLN • DML > 150% ULN • F-response latency > 120% ULN, or > 150% ULN (if distal CMAP < 50% of LLN) OR • F-wave absence in two nerves with dCMAP $\geq 20\%$ LLN, with an additional parameter, in one other nerve OR • pCMAP/dCMAP amplitude ratio < 0.7 (excluding the tibial nerve), in two nerves with an additional parameter, in one other nerve 2) Axonal GBS (including inexcitable forms) • dCMAP < 80% LLN in two nerves • F-wave absence in two nerves with distal CMAP $\geq 20\%$ LLN, in absence of any demyelinating feature in any nerve • pCMAP/dCMAP amplitude ratio < 0.7, in two nerves (excluding the tibial nerve) OR pCMAP/dCMAP amplitude ratio < 0.7, in one nerve; with IN ADDITION, dCMAP < 80% LLN in one other nerve Inexcitable • If dCMAP absent in all nerves (or present in only one nerve with dCMAP < 10% LLN) 3) Equivoal • Abnormal range findings however not fitting criteria for any other group

Uncini's criteria

1) Acute inflammatory demyelinating polyneuropathy (AIDP)
 At first or second study, at least one of the following in at least two nerves:
 • MCV < 70% LLN
 • DML > 130% ULN
 • dCMAP duration > 120% ULN
 • pCMAP/dCMAP duration > 130%
 • F-response latency > 120% ULN
 OR one of the above in one nerve PLUS:
 • Absent F waves in two nerves with dCMAP > 20% LLN
 • Abnormal ulnar SNAP amplitude and normal sural SNAP amplitude

2) Axonal GBS
 Acute motor axonal neuropathy (AMAN)
 At first and second study, none of the above AIDP features in any nerve (demyelinating features allowed in one nerve if dCMAP < 20% LLN)
 At first study, at least one of the following in each of two nerves:
 • dCMAP < 80% LLN
 • pCMAP/dCMAP amplitude ratio < 0.7 (excluding tibial nerve)
 • isolated F wave absence (or < 20% persistence)
 At second study:
 • at least one of the followings in two nerves is evidence of axonal degeneration:
 • persistent or further reduction of dCMAP amplitude
 • pCMAP/dCMAP amplitude ratio < 0.7 at first test which recovers because of decrease of dCMAP without increased temporal dispersion (dCMAP duration $\leq 120\%$ ULN and pCMAP/dCMAP duration ratio $\leq 130\%$)
 • at least one of the followings in two nerves is evidence of reversible conduction failure:
 • > 150% increase dCMAP amplitude without increased dCMAP duration ($\leq 120\%$ ULN)
 • pCMAP/dCMAP amplitude ratio < 0.7 at first test which improves more than 0.2 because of increased pCMAP without temporal dispersion (pCMAP/dCMAP duration ratio $\leq 130\%$)
 • isolated F wave absence (or < 20% persistence) that recovers without increased minimal latency ($\leq 120\%$ ULN)
 Acute motor and sensory axonal neuropathy (AMSAN)
 At first study:
 • the same criteria of AMAN in motor nerves PLUS
 • SNAP amplitudes < 50% LLN in at least two nerves
 At second study:
 • evidence for axonal degeneration and reversible conduction failure in motor nerves as in AMAN
 • there is evidence of axonal degeneration in sensory nerves if SNAP amplitude in two nerves it is stable or decreased
 • there is evidence of reversible conduction failure in sensory nerves if SNAP amplitude in two nerves it is increased (> 50% in median and ulnar nerves and > 60% in sural)
3) Inexcitable
 At first or second study:
 • Distal CMAP absent in all nerves (or present in only one with distal CMAP < 10% LLN)
4) Equivoal
 At first or second study:
 • Abnormal findings not fulfilling any of the above criteria

Guillain-Barré syndrome subtypes: A clinical electrophysiological study of 100 patients

A.-M. Grapperon
 Rev Neurol (Paris) 2019 Jan - Feb;175(1-2):73-80

	Hadden criteria	Rajabally criteria
All patients (n = 100)		
Axonal	9	41
Demyelinating	52	41
Normal	8	8
Equivoal	31	10
Classic GBS (n = 69)		
Axonal	6	29
Demyelinating	38	33
Normal	1	0
Equivoal	24	7
Miller-Fisher syndrome (MFS) (n = 8)		
Axonal	0	1
Demyelinating	2	1
Normal	6	6
Equivoal	0	0
Pharyngeal-cervical-brachial (PCB) variant (n = 2)		
Axonal	1	1
Demyelinating	0	0
Normal	0	0
Equivoal	1	1
Paraparetic Guillain-Barré syndrome (p-GBS) (n = 12)		
Axonal	4	8
Demyelinating	6	2
Normal	0	0
Equivoal	2	2
Unclassified (n = 9)		
Axonal	1	3
Demyelinating	6	1
Normal	1	1
Equivoal	1	4

Safety and efficacy of eculizumab in Guillain-Barré syndrome: a multicentre, double-blind, randomised phase 2 trial

Sonoko Misawa, Satoshi Kuwabara
Lancet Neurol 2018; 17: 519-29

	Eculizumab (n=23)	Placebo (n=11)
Age (years)	56 (42-74)	59 (32-73)
Median (IQR)		
Sex		
Men	16 (70%)	8 (73%)
Women	7 (30%)	3 (27%)
Antecedent illness		
None	2 (9%)	1 (9%)
Respiratory tract infection	11 (48%)	3 (27%)
Diarrhoea	12 (52%)	8 (73%)
Other	4 (17%)	2 (18%)
Days from onset to immunoglobulin administration		
Mean (range, SD)	5.0 (2.0-12.0, 2.6)	4.6 (2.0-13.0, 3.4)
Functional grade		
3 (able to walk 5 m with a walker or support)	3 (13%)	1 (9%)
4 (bed-bound or chair-bound)	19 (83%)	9 (82%)
5 (requires assisted respiration)	1 (4%)	1 (9%)
Sum scores of manual muscle testing		
Mean (SD)	41.5 (15.4)	34.0 (17.9)
Grip strength (average of both sides)		
Mean (SD)	7.21 (7.34)	3.03 (4.78)
Antiganglioside antibody		
GM1	6 (26%)	5 (45%)
Electrodiagnostic subtype		
Acute motor axonal neuropathy	13 (57%)	4 (36%)
Acute inflammatory demyelinating neuropathy	7 (30%)	7 (64%)
Unclassified	3 (13%)	0 (0%)

	Eculizumab (n=23)	Placebo (n=11)	Between-group difference (95% CI)	p value
Primary outcome				
Able to walk 5 m independently (functional grade ≥ 2)				
Week 4, % (90% CI)	61% (42 to 78)	45% (20 to 73)	15% (-20 to 51)	..
Number of patients	14/23	5/11	..	..
Adverse events affecting $\geq 10\%$ patients in either group				
Any event				
	Eculizumab (n=23)	Placebo (n=11)		
Insomnia	6 (26%)	1 (9%)		
Headache	4 (17%)	2 (18%)		
Nasopharyngitis	4 (17%)	0		
Constipation	4 (17%)	2 (18%)		
Rash	4 (17%)	0		
Myalgia	3 (13%)	1 (9%)		
Nausea	3 (13%)	1 (9%)		
Dyshidrotic eczema	3 (13%)	0		
Oral mucositis	3 (13%)	0		
Hepatic function abnormal	4 (17%)	2 (18%)		
Neutrophil count decrease	3 (13%)	1 (9%)		
Hyponatraemia	3 (13%)	0		
Haematuria	2 (9%)	2 (18%)		
Serious adverse events	2 (9%)	1 (9%)		
Anaphylaxis	1 (4%)	0		
Intracranial haemorrhage	1 (4%)*	0		
Brain abscess	1 (4%)*	0		
Depression	0	1 (9%)

Patient	Age (y)	Sex	EMG	CSF protein (mg/dL)	Progression after treatment	Mechanical ventilation (d)	Hospital stay (d)	Hughes grade on admission	Hughes grade on discharge
1	16	M	AMAN	21	No	5	13	5	0
2	12	M	AMAN	48	No	13	25	5	2
3	9	M	AMAN	38	No	6	20	5	1
4	9	F	AMAN	55	No	7	12	5	1
5	8	F	AMAN	60	No	5	17	5	0
6	6	F	AMAN	50	No	5	10	5	1
7	15	F	AMAN	54	No	14	30	5	1
8	10	F	AMAN	60	No	9	23	5	2
9	11	F	AMAN	29	No	6	12	5	0

Comparison of Patients' Outcome of the Previous Treatment Modalities With Zipper Strategy.

	PE ^a	IVIG ^b	PE+IVIG ^a	Zipper strategy
Mean days to stop ventilation	29	26	18	7
Mean days to unaided walking	49	51	40	24
Mean days to hospital discharge	63	53	51	18

3 Chronic Inflammatory Demyelinating Polyneuropathy

Meta-analysis of clinical features between NF155-positive and -negative CIDP patients

	OR (95% CI)	p value	p for heterogeneity	p value for bias		Number of studies
				Begg's test	Egger's test	
Subacute	2.17 (0.98, 4.84)	0.06	0.33	1.00	-	2
Cerebellar ataxia	5.69 (1.60, 20.26)	0.007	0.08	1.00	-	2
Sensory ataxia	10.79 (5.24, 22.22)	<0.001	0.23	1.00	-	2
Tremor	6.71 (3.37, 13.39)	<0.001	0.60	0.296	0.064	3
Brain lesions	2.65 (0.95, 7.37)	0.06	0.11	1.00	0.801	3
Treatment (IVIg) good response	0.12 (0.05, 0.29)	<0.001	0.12	1.00	-	2
Sex incidence (female)	0.62 (0.34, 1.13)	0.12	0.38	0.308	0.505	4

	Fingolimod 0.5 mg (n=54)	Placebo (n=52)	p value
Change in grip strength from baseline to month 6 or end of study* (kPa)			
Dominant hand			
Patients with data	53	50	-
Mean (SD)	-4.1 (20.0)	-4.8 (15.0)	-
Median (range)	-1.3 (-60.3 to 36.0)	-4.3 (-43.7 to 34.7)	-
Least squares mean difference (95% CI)†	-	1.2 (-5.9 to 8.2)	0.73
Non-dominant hand			
Patients with data	53	50	-
Mean (SD)	-4.7 (19.9)	-7.4 (14.8)	-
Median (range)	-1.0 (-55.3 to 38.3)	-4.7 (-52.7 to 34.0)	-
Least squares mean difference (95% CI)†	-	3.4 (-3.6 to 10.4)	0.33
Change in R-ODS from baseline to month 6 or end of study*			
Patients with data	54	51	-
Mean (SD)	-6.5 (11.7)	-5.8 (10.2)	-
Median (range)	-3.0 (-35.0 to 9.0)	-3.0 (-49.0 to 10.0)	-
Least squares mean difference (95% CI)†	-	-0.8 (-5.0 to 3.4)	0.70

4

Inflammatory myopathy

Muscular and extramuscular features of myositis patients with anti-U1-RNP autoantibodies

Maria Casal-Dominguez
Neurology® 2019;92:e1-e11.

Muscle biopsy features of anti-U1-ribonucleoprotein (RNP) and control groups

	Anti-U1-RNP (n = 7)	DM (n = 45)	AS (n = 27)	IMNM (n = 70)
Necrotizing myopathy	57 (4)	11 (5) ^a	22 (6)	80 (56)
Perifascicular atrophy	0 (0)	56 (25) ^a	52 (14) ^a	0 (0)
Perivascular inflammation	43 (3)	62 (28)	63 (17)	27 (19)
Primary inflammation	0 (0)	9 (4)	30 (8)	16 (11)

Cumulative clinical features of anti-U1-ribonucleoprotein (RNP)-positive patients and control groups

	Anti-U1-RNP (n = 26)	DM (n = 178)	AS (n = 132)	IMNM (n = 135)
Muscle involvement				
Muscle weakness	80 (16)	85 (152)	90 (119)	96 (130) ^a
Myalgia	60 (12)	56 (100)	65 (86)	52 (70)
Skin involvement				
DM-specific skin involvement	60 (12)	96 (171) ^a	60 (79)	4 (5) ^a
Sicca-specific skin involvement	25 (5)	2 (3) ^a	13 (17)	0 (0) ^a
Raynaud phenomenon	80 (16)	22 (40) ^a	39 (52) ^a	15 (20) ^a
Telangiectasias	20 (4)	21 (37)	20 (26)	8 (11)
Ulcers	15 (3)	14 (25)	7 (9)	0 (0) ^a
Carpal tunnel	15 (3)	8 (15)	20 (27)	10 (13)
Livedo reticularis	20 (4)	12 (22)	10 (13)	4 (5) ^a
Mechanics hands	50 (10)	28 (49) ^a	58 (77)	5 (7) ^a
Calcinosis	25 (5)	21 (38)	9 (12)	1 (1) ^a
Subcutaneous edema	35 (7)	18 (32)	27 (35)	4 (5) ^a
Puffy hands	20 (4)	8 (15)	10 (13)	0 (0) ^a
Lung involvement				
Interstitial lung disease	45 (9)	13 (24) ^a	80 (106) ^a	6 (8) ^a
Pulmonary hypertension	25 (5)	3 (5) ^a	20 (27)	1 (2) ^a
Esophageal involvement				
Gastroesophageal reflux	45 (9)	29 (51)	29 (38)	25 (34)
Dysphagia	50 (10)	53 (95)	18 (24) ^a	39 (53)
Joint involvement				
Arthritis	60 (12)	18 (32) ^a	55 (72)	6 (8) ^a
Arthralgia	65 (13)	51 (90)	62 (82)	36 (49) ^a
Systemic involvement				
Fever	35 (7)	18 (32)	24 (32)	7 (10) ^a
Sicca syndrome	15 (3)	31 (55)	48 (63) ^a	19 (26)
Pericarditis	40 (8)	0 (0) ^a	1 (1) ^a	0 (0) ^a
Glomerulonephritis	25 (5)	0 (0) ^a	1 (1) ^a	0 (0) ^a